

LATTICE COMPUTATION OF THE EFFECTIVE POTENTIAL IN O(2)-INVARIANT $\lambda\Phi^4$ THEORY

A. Agodi, G. Andronico and M. Consoli

Dipartimento di Fisica - Universita di Catania;

Istituto Nazionale di Fisica Nucleare - Sezione di Catania;

Corso Italia, 57 - I 95129 Catania - Italy

Abstract

We present a lattice computation of the effective potential for O(2)-invariant $(\lambda\Phi^4)_4$ theory in the region of bare parameters corresponding to a classically scale-invariant theory. As expected from “triviality” and as in the one-component theory, we find very good agreement with the one-loop prediction, while a perturbative leading-log improvement of the effective potential fails to reproduce the Monte Carlo data. The mass m_h of the free shifted radial field is related to the renormalized vacuum expectation value v_R through the same relation $m_h^2 = 8\pi^2 v_R^2$ as in the one-component case. This confirms the prediction of a weakly interacting 2.2 TeV Higgs particle in the standard model.

In the study of Spontaneous Symmetry Breaking (SSB) the simplest quantity to compute is the vacuum expectation value of the bare scalar field $\Phi_B(x)$ in the presence of an external constant source J

$$\langle \Phi_B \rangle_J = \phi_B(J) \quad (1)$$

Computing ϕ_B on the lattice at several J -values is equivalent [1] to inverting the relation

$$J = J(\phi_B) = \frac{dV_{eff}}{d\phi_B} \quad (2)$$

involving the effective potential $V_{eff}(\phi_B)$. In this framework, the occurrence of SSB is determined from exploring (for $J \neq 0$) the properties of the function

$$\phi_B(J) = -\phi_B(-J) \quad (3)$$

in connection with the limiting behaviour at zero external source

$$\lim_{J \rightarrow 0^\pm} \phi_B(J) = \pm v_B \neq 0 \quad (4)$$

over a suitable range of the bare parameters (r_o, λ_o) in the lattice action

$$a^4 \sum_x \left[\frac{\sum_{\mu=1}^4 (\Phi_B(x + ae_\mu) - \Phi_B(x))^2}{2a^2} + \frac{1}{2} r_o \Phi_B^2(x) + \frac{\lambda_o}{4} \Phi_B^4(x) \right] \quad (5)$$

A lattice simulation of the weakly coupled one-component massless $\lambda\Phi^4$ theory [2] has shown that there is a well defined region in the bare parameter plane (r_o, λ_o) , corresponding to SSB from the classically scale-invariant case, where the effective potential is reproduced by its one-loop form to very high accuracy. Even though $\frac{\lambda_o}{\pi^2} \ll 1$, a “leading-log improvement” of the one-loop potential completely fails to reproduce the Monte Carlo data. This result, while contradicting the naive perturbative expectations, confirms the crucial insight of [3,4] on the basis of the generally accepted “triviality” [5] of $(\lambda\Phi^4)_4$: for a “trivial” theory the one-loop potential is effectively exact in the continuum limit. In fact, since there are no observable interactions, the effective potential is just the sum of the classical potential and the zero-point energy of the free fluctuation field $h(x) = \Phi_B(x) - \langle \Phi_B \rangle$. The traditional perturbative renormalization, based on the concept of a cutoff-independent and *non-vanishing* renormalized coupling at non-zero external momenta λ_R , is not appropriate [6] just because it would spoil this exactness—it does not properly re-absorb infinities but merely pushes them into “higher orders” which are then neglected. In this sense, the unphysical features of the

perturbative renormalization (the one-loop Landau pole or the spurious two-loop ultraviolet fixed point at non zero bare coupling which contradicts the rigorous arguments of [5]) are just a signal of the inadequacy of the procedure. However, it is simple to renormalize the one-loop potential exactly [3,4,7–11] (this was first discovered in the context of the Gaussian effective potential [12–14]). The particle mass m_h is related to the ultraviolet cutoff and to the bare vacuum field v_B through

$$m_h^2 = 3\lambda_o v_B^2 = \Lambda^2 \exp\left(-\frac{16\pi^2}{9\lambda_o}\right). \quad (6)$$

[Note that $\lambda_o = \lambda_B/6$ in the notation of [3,4]]. At the same time, the vacuum energy density (a renormalization-group-invariant quantity) is

$$V^{1\text{-loop}}(\pm v_B) = -\frac{m_h^4}{128\pi^2} \quad (7)$$

Hence, to get a cutoff-independent m_h in the continuum limit $\Lambda \rightarrow \infty$, $\lambda_o = \lambda_o(\Lambda)$ has to vanish as $1/(\ln \frac{\Lambda}{m_h})$ and the bare vacuum field ϕ_B is non-trivially rescaled with respect to the physical field ϕ_R through

$$\phi_B^2 = Z_\phi \phi_R^2 \quad (8)$$

with $Z_\phi \sim 1/\lambda_o$. The non-perturbative nature of the vacuum field renormalization ($Z_\phi \sim 1/\lambda_o$), first discovered in the gaussian approximation by Stevenson and Tarrach [13], should not be confused with the h -field wave function renormalization. Since h is just a free field, one has trivially $Z_h = 1$. The structure $Z_\phi \neq Z_h$ is allowed because for a scalar field the decomposition into $p_\mu = 0$ and $p_\mu \neq 0$ components is Lorentz-invariant [3,4,11]. This structure is more general than in perturbation theory and is the essential ingredient that allows SSB to coexist with “triviality”. Finally, the physical normalization condition [8,10,3,4]

$$\frac{d^2 V^{1\text{-loop}}}{d\phi_R^2} \Big|_{\phi_R=\pm v_R} = m_h^2 \quad (9)$$

determines

$$Z_\phi = \frac{8\pi^2}{3\lambda_o} \quad (10)$$

and leads to

$$m_h^2 = 8\pi^2 v_R^2 \quad (11)$$

In the context of the standard model, where v_R is known from the Fermi constant to be 246 GeV, this predicts a Higgs mass ~ 2.2 TeV (up to radiative corrections that are small if the top mass is below 200 GeV [8,10]).

As discussed in [8,10,3], one expects Eq.(11), obtained in the single-component theory, to be also valid in the $O(N)$ -continuous symmetry case. This observation originates in Ref. [15] which obtained the same effective potential for the radial field as in the one-component theory. This is extremely intuitive. The Goldstone-boson fields do not contribute non-trivially to the effective potential: they contribute only their zero-point energy, which is a constant since these are free, massless fields, according to “triviality”. Thus, in the $O(2)$ case, one may take the diagram (V_{eff}, ϕ_B) for the one-component theory and “rotate” it around the V_{eff} symmetry axis. This generates a three- dimensional diagram $(V_{eff}, \phi_1, \phi_2)$ where V_{eff} depends on the bare radial field,

$$\rho_B = \sqrt{\phi_1^2 + \phi_2^2} \quad (12)$$

in exactly the same way as V_{eff} depends on ϕ_B in the one-component theory; namely $(\omega^2(\rho_B) = 3\lambda_o\rho_B^2)$

$$V^{1-loop}(\rho_B) = \frac{\lambda_o}{4}\rho_B^4 + \frac{\omega^4(\rho_B)}{64\pi^2} \left(\ln \frac{\omega^2(\rho_B)}{\Lambda^2} - \frac{1}{2} \right). \quad (13)$$

This represents the classical potential plus the zero-point energy of the *free* shifted radial field. “Triviality” implies that there are no observable interaction effects, so this result should be exact [3,4]. By using eqs.(6,10), V^{1-loop} can be re-expressed in the form

$$V^{1-loop}(\rho_B) = \frac{\pi^2 \rho_B^4}{Z_\phi^2} \left(\ln \frac{\rho_B^2}{v_B^2} - \frac{1}{2} \right), \quad (14)$$

Differentiating Eq.(14), we obtain the bare “radial source”

$$J(\rho_B) = \frac{dV^{1\text{-loop}}(\rho_B)}{d\rho_B} = \frac{4\pi^2 \rho_B^3}{Z_\phi^2} \ln \frac{\rho_B^2}{v_B^2}, \quad (15)$$

which we shall compare with the lattice results for $J = J(\rho_B)$.

The lattice simulation of the O(2)-invariant theory is obtained from the action

$$\begin{aligned} a^4 \sum_x & \left[\frac{\sum_{\mu=1}^4 (\Phi_1(x + ae_\mu) - \Phi_1(x))^2}{2a^2} + \frac{\sum_{\mu=1}^4 (\Phi_2(x + ae_\mu) - \Phi_2(x))^2}{2a^2} \right. \\ & \left. + \frac{1}{2} r_o (\Phi_1^2(x) + \Phi_2^2(x)) + \frac{\lambda_o}{4} (\Phi_1^2(x) + \Phi_2^2(x))^2 \right] \end{aligned} \quad (16)$$

and one couples Φ_1 and Φ_2 to two constant external sources J_1 and J_2 through

$$a^4 \sum_x [J_1 \Phi_1(x) + J_2 \Phi_2(x)] \quad (17)$$

By using $J_1 = J \cos \theta$ and $J_2 = J \sin \theta$ it is straightforward to show that the bare radial field ($\phi_1 = \langle \Phi_1 \rangle_{J_1, J_2}$, $\phi_2 = \langle \Phi_2 \rangle_{J_1, J_2}$)

$$\rho_B = \sqrt{\phi_1^2 + \phi_2^2} \quad (18)$$

does only depend on J , that is

$$\rho_B = \rho_B(J) \quad (19)$$

We started our Monte Carlo simulation on a 10^4 lattice by investigating first the (r_o, λ_o) correlation which corresponds to the classically scale-invariant case. Analytically, this corresponds to determine r_o from the zero-mass renormalization condition [16]

$$\frac{d^2 V_{eff}}{d\rho_B^2} \Big|_{\rho_B=0} = 0 \quad (20)$$

so that the theory does not contain any intrinsic scale in its symmetric phase $\langle \Phi_1 \rangle = \langle \Phi_2 \rangle = 0$. However, on the lattice, due to the relatively large errors introduced from the direct use of Eq.(20), it is more convenient to define the massless theory as in [2] for the one-component theory. There, we started from the general expression [4]

$$J(\phi_B) = \alpha \phi_B^3 \ln(\phi_B^2/v_B^2) + \beta v_B^2 \phi_B (1 - \phi_B^2/v_B^2), \quad (21)$$

which is still consistent with “triviality” (corresponding to an effective potential given by the sum of a classical background and the zero point energy of a free field) but allows for an explicit scale-breaking term β . Setting $\alpha = 0$ one obtains a good description of the data in the “extreme double well” limit (r_o much more negative than r_c , where r_c corresponds to the onset of SSB) where SSB is a semi-classical phenomenon and the zero-point energy represents a small perturbation. Then, we started to increase r_o , at fixed λ_o , toward the unknown value r_c and examined the quality of the fit with (α, β, v_B) as free parameters. The value of r_o at which the quality of the 2- parameter fit ($\alpha, \beta = 0, v_B$) becomes exactly the same as that of the more general 3- parameter case was used to define the massless case. At $\lambda_o = 1$ the massless regime of the one-component theory was found at a value $r_o = r_s$ where $r_s a^2 \sim -0.45$. Finally, the accurate weak-coupling relation between the bare mass and the euclidean cutoff [16] ($\lambda_B = 6\lambda_o$)

$$r_s = -\frac{\lambda_B}{32\pi^2} \Lambda^2 = -\frac{3\lambda_o}{16\pi^2} \Lambda^2 \quad (22)$$

was used to generate the massless theory at different values of λ_o from the previously determined value of r_s at $\lambda_o = 1$. It should be noted that our numerical coefficient relating ultraviolet cutoff and lattice spacing for quadratic divergences ($\Lambda a \sim 4.87$) agrees well with an independent analysis of lattice data presented by Brahm [17] which gives (in the range $\lambda_o \leq 10$) $\Lambda a = 4.893 \pm 0.003$. Also, ref. [17] predicts the massless regime to correspond to $r_s a^2 = -(0.224 \pm 0.001)$ for $\lambda_o = 0.5$ in the infinite-volume limit. This implies $r_s a^2 \sim -(0.448 \pm 0.002)$ for $\lambda_o = 1$, in excellent agreement with our result.

In the O(2) case, one can use the results of [2,17] by modifying Eq.(22) for the two-component case. This simply introduces a combinatorial factor of 4/3 so that Eq.(22) becomes

$$r_s = -\frac{\lambda_o}{4\pi^2} \Lambda^2 \quad (23)$$

Hence, we expect the massless case to correspond to $r_s a^2 \sim -0.6$ for $\lambda_o = 1$. This was

confirmed to good accuracy by using the above-described fitting procedure to Eq.(21) (now with ϕ_B replaced by ρ_B). Thus, the identification of the massless regime on the lattice does seem to obey the simple scaling laws (22-23) and is under theoretical control.

In the analysis of the one-component theory it was found [2] that Eq.(3) was poorly reproduced numerically at small values of J ($a^3|J| \sim 0.01$ or smaller). As a consequence, the values of ϕ_B (and any higher-order Green's functions) extracted from the direct computation at $J = 0$ were not reliable. Similarly, in the $O(2)$ case we find that at small J the exact θ -independence of ρ_B (see Eq.(19)), is poorly reproduced and our data processing becomes unreliable. We therefore consider a "safe" region of J -values, $Ja^3 \geq 0.05$, in which the spurious θ -dependence is less than $\pm 3\%$. Fitting the data to Eq.(15) we can infer the values of v_B and Z_ϕ and compare with Eqs.(6,10).

Our numerical values for $a\rho_B(J)$ in the massless case, obtained with the Metropolis algorithm on a 10^4 lattice, are reported in Table I for $\lambda_o = 1.0, 1.5$ and 2.0 . For each λ_o the corresponding r_o is computed by using Eq.(23) and our numerical input $r_s a^2 = -0.6$ for $\lambda_o = 1$. The errors in Table I are essentially determined from the observed spurious variation of ρ_B in the range $0 \leq \theta \leq 2\pi$. As discussed above, this is a numerical artifact and should be considered a systematic effect of the Monte Carlo lattice simulation. It is reproduced with three random number generators consistent with the Kolmogorov-Smirnov test at the level $O(10^{-4})$. At low J this systematic effect completely dominates the error; the statistical errors, after 30,000 iterations, are 4-5 times smaller [18].

Table I also reports the v_B and Z_ϕ values obtained from the two-parameter fits to the data using Eq.(15). The resulting Z_ϕ values agree well with the one-loop prediction in Eq.(10). To perform a more stringent test of the one-loop potential we next constrain Z_ϕ to its one-loop value in Eq.(10) and make a precise determination of av_B from a *one*-parameter fit to Eq.(15). This allows a meaningful comparison with the lattice version of Eq.(6)

$$(av_B)^{1-\text{loop}} = \frac{\pi y_L}{\sqrt{3\lambda_o}} \exp\left(-\frac{8\pi^2}{9\lambda_o}\right) \quad (24)$$

where we have identified the euclidean ultraviolet cutoff $\Lambda \rightarrow \frac{\pi y_L}{a}$, y_L being an *a priori*

unknown coefficient. By replacing Eq.(24) into Eq.(15) we determined y_L from the one-parameter fit to the data at $\lambda_o = 1.5$ and $r_o a^2 = -0.9$ to be $y_L = 2.44 \pm 0.03$ ($\frac{\chi^2}{d.o.f.} = \frac{3.3}{17}$). (As discussed in [2], one does not expect precisely the same numerical coefficient to govern the relation between euclidean cutoff and lattice spacing for both quadratic and logarithmic divergences, see below). In Table II we show the results of the one-parameter fits to the data at $\lambda_o = 1.0$ and 2.0 and the comparison with Eq.(24) for $y_L = 2.44 \pm 0.03$. It is apparent from Table II that the one-loop potential well reproduces the lattice data, as previously discovered in the one-component case [2].

The value of y_L obtained from the lattice simulation of the $O(2)$ massless $\lambda\Phi^4$ theory is $\sim 17\%$ larger than the value $y_L = 2.07 \pm 0.01$ obtained in the one-component case [2]. This is due to our choice of fixing the value of the bare mass $r_s a^2 = -0.6$ at $\lambda_o = 1$ with the simple combinatorial factor $4/3$ discussed above and ignoring finite size corrections to Eq.(23), see [17]. Also, in the $O(2)$ case, errors (as estimated from the spurious θ -dependence of ρ_B) are larger than in the one-component theory and, moreover, the massless regime appears not so sharply identified on the basis of the (J, ρ_B) correlation. Indeed, it is found in a narrow range of r_o -values near $r_o a^2 = -0.6$ for $\lambda_o = 1$. Choosing instead $r_s a^2 \sim -0.58$ would give $y_L \sim 2.07$.

The agreement between one-loop predictions and Monte Carlo data is not a trivial test of perturbation theory but, rather, represents a non perturbative test of “triviality”. If perturbation theory were valid then the data should agree at least as well, if not better, with the leading-log formula based on the perturbative β -function

$$J^{\text{LL}}(\rho_B) = \frac{\lambda_o \rho_B^3}{1 + \frac{5\lambda_o}{4\pi^2} \ln \frac{\pi x_{\text{LL}}}{a\rho_B}}$$

(x_{LL} denoting an adjustable parameter). However, when we fit the $\lambda_o = 1.0, 1.5$ and 2.0 data to this formula we find, respectively, for 17 degrees of freedom, $(\chi^2)_{\text{LL}} = 9, 44$ and 133 (to compare with the values $\chi^2 = 0.8, 3.3$ and 7.3 obtained from the 1-loop one-parameter fits with Eq.(15) when Z_ϕ is constrained to its value in Eq.(10)). Note that we are in a region where the “ $\lambda_o \log$ ” term is not small; it is of order unity. Thus, the good agreement between

the data and the one-loop formula is not because the higher-order corrections, expected by perturbation theory, are negligibly small; it is because they are *absent*. Without the “triviality” argument, this would be an incomprehensible miracle.

In conclusion, the good agreement between lattice simulation and eqs. (15,10,23,24) provides definite evidence that the dependence of the effective potential on the radial field in the continuous symmetry case is completely consistent with our expectations [3,4]. This confirms that Eq.(11), up to small radiative corrections due to the gauge and Yukawa couplings, controls the relation between the Higgs mass and the Fermi constant in the standard model if SSB is generated through “dimensional transmutation” from a classically scale-invariant $\lambda\Phi^4$ theory. The “triviality” structure we have checked, in which m_h is not proportional to “ λ_R ” (which vanishes), implies that the Higgs, despite of its rather large mass, is only weakly interacting and would be free if the gauge and Yukawa couplings would be turned off.

ACKNOWLEDGEMENTS

We thank P. M. Stevenson for very useful discussions and collaboration.

REFERENCES

- [1] K. Huang, E. Manousakis, and J. Polonyi, Phys. Rev. **D35**, 3187 (1987); K. Huang, Int. J. Mod. Phys. **A4**, 1037 (1989); in Proceedings of the DPF Meeting, Storrs, CT, 1988.
- [2] A. Agodi, G. Andronico, and M. Consoli, *The real test of “triviality” on the lattice*, INFN and University of Catania preprint, February 1994, submitted to Physical Review Letters (hep-th-9402071).
- [3] M. Consoli and P. M. Stevenson, *Resolution of the $\lambda\Phi^4$ Puzzle and a 2 TeV Higgs Boson*, Rice University preprint, DE-FG05-92ER40717-5, July 1993, submitted to Physical Review D. (hep-ph 9303256).
- [4] M. Consoli and P. M. Stevenson, *The Non-Trivial Effective Potential of the “Trivial” $\lambda\Phi^4$ Theory: A Lattice Test*, Rice University preprint, DE-FG05-92ER40717-9, October 1993, to appear in Zeit.für Physik C.
- [5] J. Fröhlich, Nucl. Phys. **B200**(FS4), 281 (1982); R. Fernández, J. Fröhlich, and A. D. Sokal, *Random Walks, Critical Phenomena, and Triviality in Quantum Field Theory* (Springer-Verlag, Berlin, 1992).
- [6] M. Consoli and P. M. Stevenson, *“Triviality” and the perturbative expansion in $\lambda\Phi^4$ theory*, Rice University preprint, March 1994, submitted to Physics Letters B.
- [7] V. Branchina, P. Castorina, M. Consoli and D. Zappala’, Phys. Lett. **B274**, 404 (1992).
- [8] M. Consoli, in “Gauge Theories Past and Future – in Commemoration of the 60th birthday of M. Veltman”, R. Akhoury, B. de Wit, P. van Nieuwenhuizen and H. Veltman Eds., World Scientific 1992, p. 81; M. Consoli, Phys. Lett. **B 305**, 78 (1993).
- [9] R. Ibañez-Meier and P. M. Stevenson, Phys. Lett. **B297**, 144 (1992).
- [10] V. Branchina, M. Consoli and N. M. Stivala, Zeit. Phys. **C57**, 251 (1993).
- [11] U. Ritschel, Phys. Lett. **B318**, 617 (1993).

- [12] M. Consoli and A. Ciancitto, Nucl. Phys. **B254**, 653 (1985).
- [13] P. M. Stevenson and R. Tarrach, Phys. Lett. **B176**, 436 (1986).
- [14] P. Castorina and M. Consoli, Phys. Lett. **B235**, 302 (1990); V. Branchina, P. Castorina, M. Consoli and D. Zappala', Phys. Rev. **D42**, 3587 (1990).
- [15] L. Dolan and R. Jackiw, Phys. Rev. **D9**, 3320, (1974).
- [16] S. Coleman and E. Weinberg, Phys. Rev. **D7**, 1888 (1973).
- [17] D. E. Brahm, *The lattice cutoff for $\lambda\Phi_4^4$ and $\lambda\Phi_3^6$* , CMU-HEP94-10 preprint, hep-lat/9403021, March 1994.
- [18] We have found, however, that the use of the Langevin formulation of the lattice theory considerably reduces this unphysical effect thus allowing, in principle, a more precise determination of the (J, ρ_B) correlation. We are presently investigating this alternative approach.

TABLES

TABLE I. The values of $a\rho_B(J)$ for the massless case are reported as discussed in the text. At the various values of λ_o and r_o we also show the results of the 2-parameter fits with Eq.(15) and the one loop prediction (10).

Ja^3	$\lambda_o = 1.0 \ r_o a^2 = -0.6$	$\lambda_o = 1.5 \ r_o a^2 = -0.9$	$\lambda_o = 2.0 \ r_o a^2 = -1.2$
0.050	0.4110 ± 0.0130	0.3850 ± 0.0116	0.3753 ± 0.0091
0.075	0.4686 ± 0.0086	0.4337 ± 0.0086	0.4176 ± 0.0069
0.100	0.5133 ± 0.0084	0.4724 ± 0.0065	0.4517 ± 0.0055
0.125	0.5495 ± 0.0063	0.5048 ± 0.0065	0.4814 ± 0.0054
0.150	0.5819 ± 0.0060	0.5332 ± 0.0053	0.5063 ± 0.0042
0.200	0.6377 ± 0.0047	0.5812 ± 0.0040	0.5500 ± 0.0037
0.250	0.6844 ± 0.0039	0.6217 ± 0.0034	0.5875 ± 0.0029
0.300	0.7256 ± 0.0030	0.6571 ± 0.0026	0.6190 ± 0.0029
0.350	0.7612 ± 0.0029	0.6892 ± 0.0026	0.6473 ± 0.0026
0.400	0.7938 ± 0.0025	0.7178 ± 0.0025	0.6731 ± 0.0025
0.450	0.8246 ± 0.0024	0.7435 ± 0.0025	0.6969 ± 0.0023
0.500	0.8520 ± 0.0023	0.7683 ± 0.0021	0.7193 ± 0.0023
0.550	0.8785 ± 0.0021	0.7911 ± 0.0021	0.7399 ± 0.0018
0.600	0.9029 ± 0.0019	0.8124 ± 0.0020	0.7593 ± 0.0017
0.650	0.9261 ± 0.0019	0.8330 ± 0.0019	0.7778 ± 0.0017
0.700	0.9481 ± 0.0018	0.8520 ± 0.0019	0.7953 ± 0.0017
0.750	0.9693 ± 0.0018	0.8703 ± 0.0017	0.8119 ± 0.0015
0.800	0.9893 ± 0.0016	0.8879 ± 0.0015	0.8274 ± 0.0013
	$Z_\phi = 25.0 \pm 1.6$	$Z_\phi = 16.4 \pm 0.7$	$Z_\phi = 12.3 \pm 0.4$
	$av_B = (1.4^{+1.5}_{-0.9})10^{-3}$	$av_B = (1.8 \pm 0.6)10^{-2}$	$av_B = (5.6 \pm 0.8)10^{-2}$
	$Z_\phi^{1-\text{loop}} = 26.3$	$Z_\phi^{1-\text{loop}} = 17.5$	$Z_\phi^{1-\text{loop}} = 13.1$

TABLE II. By using Eq.(15), we show the results of the 1-parameter fits for av_B at $\lambda_o = 1.0$ and 2.0 when Z_ϕ is constrained to its one-loop value in Eq.(10). We also show the predictions from Eq.(24), $(av_B)^{\text{Th}}$, for $y_L = 2.44 \pm 0.03$ as determined from the fit to the data at $\lambda_o = 1.5$.

$\lambda_o = 1.0 \quad r_o a^2 = -0.6$	$\lambda_o = 2.0 \quad r_o a^2 = -1.2$
$Z_\phi = 26.32 = \text{fixed}$	$Z_\phi = 13.16 = \text{fixed}$
$av_B = (7.09 \pm 0.11)10^{-4}$	$av_B = (3.79 \pm 0.03)10^{-2}$
$\frac{\chi^2}{d.o.f} = \frac{0.8}{17}$	$\frac{\chi^2}{d.o.f} = \frac{7.3}{17}$
$(av_B)^{\text{Th}} = (6.85 \pm 0.09)10^{-4}$	$(av_B)^{\text{Th}} = (3.89 \pm 0.05)10^{-2}$